



Review

# Cyanogenic glucosides and plant–insect interactions

Mika Zagrobelny<sup>a</sup>, Søren Bak<sup>a</sup>, Anne Vinther Rasmussen<sup>a</sup>,  
Bodil Jørgensen<sup>b</sup>, Clas M. Naumann<sup>c</sup>, Birger Lindberg Møller<sup>a,\*</sup>

<sup>a</sup>*Plant Biochemistry Laboratory, Department of Plant Biology and Center of Molecular Plant Physiology (PlaCe), Royal Veterinary and Agricultural University, 40 Thorvaldsensvej, DK-1871 Frederiksberg C, Copenhagen, Denmark*

<sup>b</sup>*Biotechnology Group, Danish Institute of Agricultural Sciences, Royal Veterinary and Agricultural University, 40 Thorvaldsensvej, DK-1871 Frederiksberg C, Copenhagen, Denmark*

<sup>c</sup>*Alexander Koenig Research Institute and Museum of Zoology, Leibniz Institute for Research in Terrestrial Biodiversity, 160 Adenauerallee, D-53113 Bonn, Germany*

Received 23 September 2003; received in revised form 9 October 2003

## Abstract

Cyanogenic glucosides are phytoanticipins known to be present in more than 2500 plant species. They are considered to have an important role in plant defense against herbivores due to bitter taste and release of toxic hydrogen cyanide upon tissue disruption. Some specialized herbivores, especially insects, preferentially feed on cyanogenic plants. Such herbivores have acquired the ability to metabolize cyanogenic glucosides or to sequester them for use in their predator defense. A few species of Arthropoda (within Diplopoda, Chilopoda, Insecta) are able to de novo synthesize cyanogenic glucosides and, in addition, some of these species are able to sequester cyanogenic glucosides from their host plant (Zygaenidae). Evolutionary aspects of these unique plant–insect interactions with focus on the enzyme systems involved in synthesis and degradation of cyanogenic glucosides are discussed.

© 2003 Elsevier Ltd. All rights reserved.

**Keywords:** Cyanogenic glucosides; Cyanogenesis; Linamarin; Lotaustralin; Lepidoptera; Zygaenidae; Papilionoidea

## Contents

1. Introduction .....	294
2. Biosynthesis, degradation and detoxification of cyanogenic glucosides .....	295
3. Cyanogenic glucosides and plant–herbivore interactions .....	296
4. Cyanogenic glucosides in Arthropoda .....	297
5. Cyanogenic glucosides in Zygaenidae (foresters and burnets) .....	298
5.1. Linamarin and lotaustralin distribution in Zygaenidae .....	298
5.2. Defensive secretion and cuticular cavities .....	299
5.3. Metabolism, catabolism and detoxification of cyanogenic glucosides .....	300
6. Cyanogenic glucosides in Papilionoidea (butterflies) .....	301
6.1. Linamarin and lotaustralin distribution .....	301
6.2. Detoxification, biosynthesis and sequestration of cyanogenic glucosides .....	301

\* Corresponding author: Tel.: +45-352-833-52; fax.: +45-352-833-33.  
E-mail address: [blm@kvl.dk](mailto:blm@kvl.dk) (B.L. Møller).

7. Conclusions and perspectives.....	302
Acknowledgements.....	303
References .....	303

## 1. Introduction

Cyanogenic glucosides (CNGs) are phytoanticipins widely distributed in the plant kingdom (Conn, 1980; Møller and Seigler, 1999; Poulton, 1990). They are present in more than 2500 different plant species including ferns, gymnosperms and angiosperms. This indicates that the ability of plants to produce CNGs is ancient. In addition, CNGs have been found in a few arthropod clades. CNGs are  $\beta$ -glucosides of  $\alpha$ -hydroxynitriles derived from the aliphatic protein amino acids L-valine, L-isoleucine and L-leucine, from the aromatic amino acids L-phenylalanine and L-tyrosine and from the aliphatic non-protein amino acid cyclopentenyl-glycine (Fig. 1). In plants, CNGs are stored in the vacuoles (Vetter, 2000). When plant tissue is disrupted e.g. by herbivore attack, CNGs are brought into contact with  $\beta$ -glucosidases and  $\alpha$ -hydroxynitrile lyases that hydrolyze the CNGs and thereby cause release of toxic hydrogen cyanide (HCN) (Fig. 1). This binary system—

two sets of components that, when separated, are chemically inert—provides plants with an immediate defense against intruding herbivores and pathogens that cause tissue damage.

Cyanide is a toxic substance, mainly due to its affinity for the terminal cytochrome oxidase in the mitochondrial respiratory pathway (Brattsten et al., 1983). The lethal dose of cyanide for vertebrates lies in the range of 35–150  $\mu\text{mol kg}^{-1}$ , if applied in a single dose. Much higher amounts of HCN can be tolerated if consumed or administered over a longer period (Davis and Nahrstedt, 1985). Biosynthesis and degradation of CNGs are well documented in many plants (Jones et al., 2000; Lechtenberg and Nahrstedt, 1999).

For most plants it has been hypothesized that CNGs are involved in plant defense against herbivores due to release of toxic HCN (Nahrstedt, 1996). CNGs are, however, also known to act as both feeding deterrents and phagostimulants for herbivores that are specialists on plants containing CNGs (reviewed in Gleadow and




Fig. 1. Biosynthesis, catabolism and detoxification of CNGs in plants, insects and higher animals. Enzymes involved are shown in red. HCN is highlighted in purple.

Woodrow, 2002). This review will aim to summarize current knowledge on CNGs and their synthesis and degradation in insects compared to plants, and to discuss possible evolutionary implications of these relationships.

## 2. Biosynthesis, degradation and detoxification of cyanogenic glucosides

The main metabolic processes resulting in synthesis, degradation and detoxification of CNGs in plants are shown in Fig. 1. The first two committed steps in CNG biosynthesis are catalyzed by cytochromes P450 (Fig. 1). The first P450 catalyzed step proceeds via two successive *N*-hydroxylations of the amino group of the parent amino acid, followed by decarboxylation and dehydration (Sibbesen et al., 1994). The aldoxime formed is subsequently converted to an  $\alpha$ -hydroxynitrile through the action of a second cytochrome P450 (Bak et al., 1998; Kahn et al., 1997) (Fig. 1). This reaction involves an initial dehydration reaction that forms a nitrile and is followed by hydroxylation of the alpha carbon to generate a cyanohydrin. The final step in CNG synthesis, glycosylation of the cyanohydrin moiety, is catalyzed by a UDPG-glycosyltransferase (Jones et al., 1999) (Fig. 1). Catabolism of CNGs is initiated by enzymatic hydrolysis by a  $\beta$ -glucosidase to afford the corresponding  $\alpha$ -hydroxynitrile, which at pH values above 6 spontaneously dissociates into a sugar, a keto compound, and HCN (Fig. 1). At lower pH values, the dissociation reaction is catalyzed by an  $\alpha$ -hydroxynitrile lyase. HCN is detoxified by two main reactions (Møller and Poulton, 1993) (Fig. 1). The first route involves the formation of  $\beta$ -cyanoalanine from cysteine and is catalyzed by  $\beta$ -cyanoalanine-synthase (Fig. 1, route 1).  $\beta$ -Cyanoalanine is subsequently converted into asparagine (Miller and Conn, 1980). The second route proceeds by conversion of HCN into thiocyanate and is catalyzed by rhodanese (Bordo and Bork, 2002) (Fig. 1, route 2). The detoxification route involving  $\beta$ -cyanoalanine is common in plants and possibly also in insects, while the thiocyanate pathway occurs mainly in vertebrates but also in some plants and insects.

Cytochromes P450 are heme-thiolate monooxygenases ubiquitously found in all kingdoms. They constitute a large group of polyphyletic proteins with amino acid sequence similarity as low as 20% within species. In eukaryotes, cytochromes P450 are type II membrane enzymes, N-terminally anchored to the endoplasmatic membrane. In prokaryotes, cytochromes P450 are soluble (Werck-Reichhart and Feyereisen, 2000). In insects and plants, cytochromes P450 are encoded by some of the largest multigene families, with 89 genes in the *Drosophila melanogaster* (Endopterygota) genome (Tijet et al., 2001) (<http://p450.antibes.inra.fr/>) (Fig. 2), 100 genes in the *Anopheles gambiae*




Fig. 2. Simplified schematic evolutionary tree for Arthropoda. Arthropod groups encompassing species that contain aromatic CNGs are shown in bold whereas the group that contains aliphatic CNGs is shown in bold and underlined. The number of species that contain CNGs within each of the groups is listed in parentheses.

(Endopterygota) genome (Ranson et al., 2002) (<http://p450.antibes.inra.fr/>) (Fig. 2) and 272 genes in the *Arabidopsis thaliana* genome (Werck-Reichhart et al., 2000) (<http://biobase.dk/P450/>). Cytochromes P450 catalyze a highly diverse range of chemical reactions that include C-hydroxylations and epoxidations, N- and S-oxidations, dehydrations and O-, N- and S-dealkylations (Feyereisen, 1999; Halkier, 1996; Morant et al., 2003).

In insects, cytochromes P450 are involved in biosynthesis of ecdysteroids and juvenile hormones as well as in metabolism and detoxification of insecticides. Cytochromes P450 play crucial roles in defense against natural products that insects have to fend off in order to be able to feed on otherwise toxic plants. The ability of an insect cytochrome P450 to metabolize a specific natural product is often the key to the adaptation of insect herbivores to their host plants (Feyereisen, 1999). In plants, cytochromes P450 are involved in biosynthesis of a vast array of natural products involved in plant defense as well as many other biosynthetic pathways (for recent reviews see Morant et al., 2003; Werck-Reichhart et al., 2002).

The final step in biosynthesis of CNGs is catalyzed by a Family 1 glycosyltransferase (Jones et al., 1999; Vogt and Jones, 2000). Family 1 glycosyltransferases are soluble proteins with a molecular mass of 45–60 kDa, which utilize UDP-activated sugar moieties as the donor molecules to glycosylate the acceptor molecules. Glycosyltransferases generally exhibit a low degree of overall sequence similarity, and are often regioselective or regiospecific rather than highly substrate specific (Hansen et al., 2003). Typically, the first committed step in a biosynthetic pathway is catalyzed by an enzyme with high substrate specificity. This serves to limit the number of available substrates for subsequent enzymes in the same pathway. These enzymes may thus possess a wider substrate specificity that provides overall

metabolic flexibility, yet desired specificity with a limited number of genes (Vogt and Jones, 2000). As for cytochromes P450, Family 1 glycosyltransferases are encoded by a multigene family and are ubiquitously found in plants, animals, fungi, bacteria and viruses (Paquette et al., 2003).

In plants, degradation of CNGs is catalyzed by  $\beta$ -glucosidases and  $\alpha$ -hydroxynitrile lyases (Conn, 1980; Hösel and Conn, 1982; Poulton, 1990).  $\beta$ -Glucosidases that catalyze the hydrolysis of glycosidic linkage in aryl and alkyl  $\beta$ -glucosides and in cellulose are present in bacteria, fungi, plants and animals. In contrast to the well characterized  $\beta$ -glucosidases involved in CNG catabolism in plants (Cicek et al., 2000; Cicek and Esen, 1998; Czjzek et al., 2000), only little is known about these insect  $\beta$ -glucosidases and their substrate specificity.  $\beta$ -Glucosidases generally have a subunit molecular mass of 55–65 kDa, acidic pH optima (pH 5–6) and an absolute specificity towards  $\beta$ -glucosides (Esen, 1993). Plant  $\beta$ -glucosidases involved in cleavage of CNGs exhibit a high specificity towards the aglycone moiety of CNGs present in the same plant species (Hösel et al., 1987; Hösel and Conn, 1982; Nahrstedt, 1985).

$\alpha$ -Hydroxynitrile lyases have been characterized in plants (Hu and Poulton, 1997, 1999; Wajant and Pfizenmaier, 1996) and in a single case from an insect (Müller and Nahrstedt, 1990). In plants, they appear to be located in the same tissues as the CNG degrading  $\beta$ -glucosidases, though their activity is observed in protein bodies (Swain et al., 1992), instead of in chloroplasts or apoplastic space as typically reported for  $\beta$ -glucosidases (Hickel et al., 1996).  $\alpha$ -Hydroxynitrile lyases constitute two broad phylogenetically distinct groups that have convergently evolved to the same function (Møller and Poulton, 1993). One homogeneous group comprises monomeric FAD-containing glycosylated enzymes. These enzymes have only been found in two subfamilies within the Rosaceae and utilize the aromatic cyanohydrin mandelonitrile as a substrate. The other group is more heterogeneous and comprises dimeric or oligomeric non-FAD containing enzymes that typically are not glycosylated. These enzymes have been found in di- and monocotyledonous plant families, and their natural substrates may be *p*-hydroxymandelonitrile as well as acetone cyanohydrin, the latter being the most common substrate. This group of non-FAD containing  $\alpha$ -hydroxynitrile lyases may be divided into subgroups depending on highest sequence similarity to serine carboxypeptidases, alcohol dehydrogenases or to rice proteins of unknown function (Hickel et al., 1996; Trummler and Wajant, 1997).

$\beta$ -Cyanoalanine synthase activity is generally found in plants (Miller and Conn, 1980) and plays a pivotal role in detoxification of HCN released as a result of cleavage of cyanogenic glucosides, or formed in stoichiometric amounts with ethylene by the enzyme 1-aminocyclo-

propane-1-carboxylic acid oxidase (Yip and Yang, 1988).  $\beta$ -Cyanoalanine synthase activity in plants and insects is primarily located in mitochondria, the organelle that is most vulnerable to HCN toxication (Meyers and Ahmad, 1991). In plants,  $\beta$ -cyanoalanine synthase has pyridoxal phosphate as a cofactor (Ikegami et al., 1988) and is a member of an ancestral family of  $\beta$ -substituted alanine synthases that also includes cysteine synthase (Ikegami and Murakoshi, 1994). Cysteine synthase also possesses  $\beta$ -cyanoalanine synthase activity and vice versa. Typically,  $\beta$ -cyanoalanine is converted into asparagine by the action of  $\beta$ -cyanoalanine hydrolase (Catric et al., 1972).  $\beta$ -Cyanoalanine is a potent neurotoxin and its accumulation in some plants may serve to deter predators (Ressler et al., 1969).

In contrast to  $\beta$ -cyanoalanine synthase, rhodanese is not ubiquitously present in plants. The *in vivo* function of rhodanese is not well understood. In those species of higher animals, plants and insects in which rhodanese is present, it is thought to play a role in cyanide detoxification (Beesley et al., 1985). In plants, this assignment is supported by high levels of rhodanese activity in 3-day-old etiolated *Sorghum bicolor* seedlings (Miller and Conn, 1980). In these seedlings, the cyanide potential is exceptionally high (Halkier and Møller, 1989). Rhodanese enzymes probably serve a variety of other functions, the most important of which is to donate sulfur to proteins (Bordo and Bork, 2002).

### 3. Cyanogenic glucosides and plant–herbivore interactions

Herbivores react very differently to the presence of CNGs in their diet. Nearly all of the variability in the effectiveness of CNGs in plant defense against herbivory is explained by four confounding factors (Gleadow and Woodrow, 2002). First, the concentration of CNGs in a host plant may be below threshold toxicity. Second, the herbivore may be a specialist that has evolved mechanisms to cope with high levels of HCN in the diet. Third, the cyanogenic plant may be consumed as part of a mixed diet and the toxicity of CNGs in this way diluted to below threshold value. Fourth, the mode of herbivore feeding may be adapted to minimize tissue damage to leaves (e.g. aphids, which are phloem feeders) to limit exposure of CNGs to degradative  $\beta$ -glucosidases (Gleadow and Woodrow, 2002). It appears that the prime deterrent effect of CNGs is linked to the keto compound released in equimolar amounts to HCN during CNG degradation, rather than to the CNG or HCN (Jones, 1988). The biosynthetic pathway for CNGs is highly channeled, preventing intermediates to dissociate from the enzyme complex (Møller and Conn, 1980; Tattersall et al., *in press*). Accordingly, the biosynthetic intermediates are not likely to act as deterrents to herbivores. In contrast, degradation of CNGs may result in accumulation of

cyanohydrins, keto compounds or aldehydes, HCN,  $\beta$ -cyanoalanine, thiocyanate and sulfite (Fig. 1). Each of these compounds may be envisioned to possess defensive properties: CNGs have a bitter taste and have been shown to act as feeding deterrents; aldehydes and ketones possess cytotoxic activities; HCN is a powerful inhibitor of respiration and of enzymes that contain heavy metals;  $\beta$ -cyanoalanine is a neurotoxin (Nahrstedt, 1985; Ressler et al., 1969) and thiocyanate and sulfite are enzyme inhibitors.

A range of plant natural products comprising alkaloids, phenylpropanoids, terpenoids, glucosinolates and CNGs have been tested for their deterrence and post-ingestional effects on *Schistocerca americana* (grasshopper, Neoptera) and *Hypera brunneipennis* (alfalfa weevil, Coleoptera) (Fig. 2). None of the compounds tested were detrimental to the grasshoppers, but eight out of ten compounds deterred feeding (Bernays, 1991; Bernays and Cornelius, 1992). Weevils were deterred by 11 of 15 compounds but again, none of these had detrimental effects. Therefore, grasshoppers and weevils seem to be behaviorally more sensitive to plant natural products, including CNGs, than required for reasonable protection from ingestion (Bernays, 1991). This may also apply to other insects that are deterred by CNGs.

The entire pathway for synthesis of the aromatic tyrosine-derived CNG dhurrin in *S. bicolor* has been transferred to *A. thaliana* using gene technology to insert the three *S. bicolor* genes *CYP79A1*, *CYP71E1* and *UGT85B1* (Tattersall et al., 2001). The accumulation of dhurrin in the transgenic *A. thaliana* plants prevented feeding by *Phyllotreta nemorum* (Coleoptera) (Fig. 2), thus unambiguously demonstrating that CNGs can confer resistance to herbivory (Tattersall et al., 2001).

In nature, some *Lotus corniculatus* plants are cyanogenic due to the presence of the two cyanogenic glucosides linamarin and lotaustralin, whereas other plants are acyanogenic either because they do not synthesize CNGs, or because they lack the  $\beta$ -glucosidase required for degradation and HCN release (Jones, 1962, 1988). Between different insects, response to the presence of CNGs in *L. corniculatus* leaves varies from total indifference to evident distaste. After starvation, insects are generally more willing to feed on cyanogenic *L. corniculatus* leaves. This indicates that for each herbivore, the deterrent capabilities of CNGs is dependent on the immediate demand for food calories (Compton and Jones, 1985). Among those species that rely on *L. corniculatus* as a major food source, there was a general lack of selectivity against CNGs, which probably reflects specialized adaptations for a cyanogenic diet. High tolerance to CNGs may be characteristic of many polyphagous Lepidoptera species, and accordingly, the role of CNGs in protection of plants from herbivores must be assessed on a species to species basis.

As the above examples imply, the primary defensive role of CNGs in plants may be as a feeding deterrent and not as a toxin (Compton and Jones, 1985). In some plants, CNGs may serve as a warning to generalist herbivores that the plant is unpalatable. CNGs are well suited to such a role, because they are recognized by a wide variety of herbivores, are a relatively cheap type of plant defense, and because HCN liberation only occurs after tissue damage, hence conserving materials and reducing the risks that adapted herbivores gain the ability to use HCN as an attractant (Compton and Jones, 1985). In conclusion, the most likely overall function of CNGs appears to be to deter herbivores that would casually try to feed on cyanogenic plants (Jones, 1988).

#### 4. Cyanogenic glucosides in Arthropoda

In contrast to the taxonomically widespread distribution of CNGs within the plant kingdom, presence of CNGs in animals appears to be restricted to a single phylum out of the currently known 31, namely the Arthropoda (Duffey, 1981; Nahrstedt, 1996) (Fig. 2). Within arthropods, presence of CNGs seems to be restricted to members of Chilopoda (centipedes), Diplopoda (millipedes) and in particular to Insecta (Davis and Nahrstedt, 1985). Within Insecta, CNGs have hitherto only been found in Coleoptera (beetles), Heteroptera (true bugs) and in particular in the Lepidoptera (butterflies and moths) (Nahrstedt, 1988) (Fig. 2).

Chilopoda, Diplopoda and some Coleoptera (*Paropsis atomaria*, *Chrysophtharta variicollis* and *C. amoena*) contain aromatic CNGs in their defensive secretions. Three beetle species appear to *de novo* synthesize CNGs as these are not present in their diet (Davis and Nahrstedt, 1985). Two species of diplopods (*Oxidus gracilis* and *Harpaghe haydeniana*) have evolved biochemical pathways for cyanogenic glucoside biosynthesis and degradation that involve very similar or identical intermediates compared to those known to be used by higher plants (Duffey, 1981). *H. haydeniana* has cyanogenic glands that contain  $\beta$ -glucosidase and  $\alpha$ -hydroxynitrile lyase activity, physically separated from the part containing CNGs. This prevents untimely release of HCN (Duffey and Towers, 1978), but offers the possibility of immediate and combined ejection and thereby mimics the phytoanticipin defense effect in plants.

In contrast to other arthropod groups, many members of the Lepidoptera are able to *de novo* synthesize CNGs as well as to sequester CNGs from their host plants (see Sections 5 and 6). Only a single species of another arthropod group, the Heteroptera, has been proposed to be able to sequester CNGs from its host plant (Braekman et al., 1982). Furthermore, members of the Lepidoptera contain mainly aliphatic CNGs as opposed to other arthropod groups that are only known to contain aromatic CNGs.

Insecta evolved at least 390 Myr ago (Gaunt and Miles, 2002), so Diplopoda and Chilopoda have evolved from a common ancestor they shared with Hexapoda at an earlier time point. This time span combined with the fact that many hexapod groups do not contain CNGs, and that those groups that do, contain different types of CNGs, points to convergent evolution of CNGs in these clades rather than to homology (Duffey, 1981). It should be emphasized that to date, the presence of CNGs has only been examined in a few species of arthropods, so the distribution of CNGs may in fact be broader than currently recognized.

### 5. Cyanogenic glucosides in Zygaenidae (foresters and burnets)

Resistance of *Zygaena* species to HCN has been well known from the beginning of the 20th century (Davis and Nahrstedt, 1985; Levinson et al., 1973). *Zygaena* species can for example remain in a concentrated atmosphere of HCN for an hour and still revive quickly when removed to clean air (Naumann et al., 1999). In 1962, the release of HCN from crushed tissues of several *Zygaena* species was shown even after the insects were reared on acyanogenic plants. This observation suggested that *Zygaena* was able to de novo synthesize CNGs (Davis and Nahrstedt, 1982; 1987; Franzl et al., 1986; Holzkamp and Nahrstedt, 1994; Jones et al., 1962). With the identification of  $\beta$ -cyanoalanine synthase in *Zygaena* larvae (Witthohn and Naumann, 1987a), it became apparent that *Zygaena* species possess the complete enzyme systems to effectively produce, control and detoxify HCN. This defense system is proposed to have evolved as a result of a strong selection pressure for protection against insectivores (Nahrstedt, 1993). The potential of Zygaenidae species to detoxify HCN as well as to de novo synthesize CNGs, is probably a basic characteristic of Zygaenidae, and may have enabled some species to commence feeding on cyanogenic plants. This would be in agreement with the observed shift of host-plant specificity from Celastraceae to Fabaceae within the Zygaenidae. This shift seems to have been very successful as evidenced by the subsequent radiation of the subgenera *Zygaena* and *Argumentia* (Zygaenidae) (Müller et al., 1993). As a special adaptation, some highly evolved *Zygaena* species, such as *Z. trifolii*, have acquired the ability to sequester CNGs from their host plants. Accordingly, these moths manage to optimize their supply of CNGs by feeding on appropriate plant sources while minimizing the energy spent to achieve this goal (Nahrstedt, 1988).

#### 5.1. Linamarin and lotaustralin distribution in Zygaenidae

The Zygaenidae are currently divided into four sub-families of which CNGs have been found in three:

Zygaeninae, Procrarinae and Chalcosiinae (Fig. 3). In addition, CNGs have been found in Charideinae and Anomoetinae, two groups that were formerly placed in the Zygaenidae, but whose positions are currently unresolved. A total of 45 species from these five groups have been shown to contain the CNGs linamarin and lotaustralin (Fig. 4) independently of the presence or absence of CNGs in the food plants provided (Davis and Nahrstedt, 1982, 1985). This indicates de novo synthesis of linamarin and lotaustralin within the Zygaeninae (Fig. 3). In agreement with previous results, we have observed the presence of linamarin and lotaustralin in several life stages of *Z. transalpiina* (Fig. 5) by LC-MS profiling (Fig. 6). In addition, both linamarin and lotaustralin were detected in the closely related families Heterogynidae, Megalopygidae and Limacodidae (Witthohn and Naumann, 1987a) (Fig. 3). Accordingly, it appears as if the accumulation of linamarin and lotaustralin is a phylogenetically old and monophyletic feature of the Zygaenoidea and perhaps also of the Zygaenoidea (Fig. 3).

Linamarin and lotaustralin appear to be sequestered from host plants (Fabaceae, e.g. *L. corniculatus*) in all examined *Zygaena* species (Nahrstedt, 1989). In




Fig. 3. Simplified schematic evolutionary tree depicting lepidopteran groups. All lepidopteran groups shown in bold and underlined contain linamarin, lotaustralin and  $\beta$ -cyanoalanine in at least one life-stage (Witthohn and Naumann, 1987a).




Fig. 4. The cyanogenic glucosides linamarin and lotaustralin derived from valine and isoleucine, respectively.




Fig. 5. *Zygaena transalpiina* life-cycle: Eggs are laid in July usually on the larval food-plant. After 1–3 weeks they hatch and the first instar larvae emerge (L1). Molting occurs every 8–10 days and the larvae stop feeding and enter diapause usually in the third, fourth or fifth instar (L3, L4, L5). During diapause, larvae reduce their metabolic turnover by about 70% compared to feeding instars and develop a considerably thicker cuticle. Larvae normally recommence feeding in spring when fresh food plants are available, but diapause may sometimes last for two or more years. The larvae reach maturity in the sixth or seventh instar (L6, L7) and 4–6 weeks after emerging from diapause, they proceed to spin cocoons and pupate. The imago moths emerge from the cocoons 14–30 days later, mate and the females lay eggs. The imagines survive 2–20 days. This life-cycle description applies to most zygaenid species (Naumann et al., 1999).

*Z. filipendulae* larvae, the amount of lotaustralin was always greater than the amount of linamarin. This probably reflects that lotaustralin is the major CNG in most leaves of one of their food plants *L. corniculatus*. A large natural variation in the linamarin and lotaustralin content in *L. corniculatus* has been reported (Gebrehiwot and Beuselinck, 2001). We confirmed this by analyzing *L. corniculatus* plants collected at three different sites in the greater Copenhagen area (Fig. 7). After pupation and in the subsequent developmental stages, linamarin was the dominant CNG in *Z. filipendulae* and also in other examined Zygaenidae (Davis and Nahrstedt, 1982). Our data obtained with *Z. transalpiina* (Fig. 6) show a more equal distribution of linamarin and lotaustralin in both larvae and imagines, although eggs contained more linamarin and empty pupae contained more lotaustralin. Accordingly, the relative amounts of linamarin and lotaustralin accumulated in Zygaenidae may vary, partly due to the relative amounts present in their diet, and partly because of the ratio generated in the insect by de novo synthesis.

In *Z. trifolii*, only small amounts of CNGs (<1%) were found in the gut and the fat body while the majority

was present in haemolymph and integument, including the defensive fluid (Davis and Nahrstedt, 1982; Franzl et al., 1986). In many insects, a large proportion of the accumulated toxic secondary plant products may be excreted or lost with exuviae during the molt. As opposed to this, *Zygaena* larvae are able to retrieve CNGs from the old cuticle (Franzl et al., 1988), since exuviae contain only minute amounts of CNGs (Fig. 6).

### 5.2. Defensive secretion and cuticular cavities

As a defensive reaction against predators (shrews, hedgehogs, starlings, frogs and carabid beetles), larvae of Zygaenini species may release highly viscous, colorless fluid droplets from cuticular cavities placed on their dorsal side (Fig. 8). Droplets appear on the cuticular surface upon contraction of irritated segments (Franzl and Naumann, 1985). The defensive fluid from *Z. trifolii* has been shown to be composed of linamarin and lotaustralin (7% CNGs) (Fig. 6),  $\beta$ -cyanoalanine (0.3%), proteins (8%, including  $\beta$ -glucosidase) and water (Witthohn and Naumann, 1984). A linamarin:lotaustralin ratio of 1:1 was measured in the defensive secretion whereas that of the haemolymph of




Fig. 6. Linamarin and lotaustralin in *Zygaena transalpiina* stages as monitored by LC–MS. Total ion traces are shown in black and overlaid with selected  $m/z$  ion traces for linamarin (green) and lotaustralin (blue). The number of specimens used for each sample is shown in parentheses. Due to a large biological variation, the signal intensity shown in the different panels should only be considered semi quantitative.




Fig. 7. Linamarin and lotaustralin polymorphism in three different populations of *Lotus corniculatus* from the larger Copenhagen area in Denmark as monitored by LC–MS. Total ion traces are shown in black, overlaid with selected  $m/z$  ion traces for linamarin (green) and lotaustralin (blue).

the *Zygaena* larvae was 19:1 (Franzl et al., 1986). This indicates that lotaustralin is transported more effectively than linamarin, maybe as a result of increased lipid solubility caused by its longer aliphatic side chain (Franzl et al., 1986). Alternatively, the shifted ratio could reflect a slower turnover rate of lotaustralin compared to linamarin in larvae.

Two morphologically different types of cavities have been found in *Z. trifolii* (Fig. 8); the larger cavities release their contents as a response to a slight irritation, whereas the smaller cavities react following severe irritation and release much smaller droplets. Defence droplets may be reabsorbed a few seconds after irritation has stopped. In contrast to most diplopods and chilopods that have specialized cyanogenic glands (Duffey, 1981), there are no gland cells or cuticular ducts leading through the cuticle into the cavities in *Zygaena* larva, and no special morphological adaptation for secretion has been developed in the epidermis (Franzl and Naumann, 1985).

### 5.3. Metabolism, catabolism and detoxification of cyanogenic glucosides

In plants, the same enzyme system uses valine and isoleucine as precursors for the synthesis of linamarin and lotaustralin, respectively, but with different catalytic efficiencies toward the substrate amino acids. This is reflected in the relative amounts of linamarin and lotaustralin in cassava (Andersen et al., 2000) and in *L. japonicus* (Forslund et al., submitted). In plants, exogenously administered *N*-hydroxyamino acids, aldoximes and nitriles can be incorporated into CNGs (Jones et al., 2000; Møller and Seigler, 1999). These same results are obtained with *Zygaena* species, and suggests that the biosynthetic pathway for CNGs in *Zygaena* is




Fig. 8. Last instar larvae of *Zygaena transalpiina* with enlarged defence droplet (top panel). Cuticular cavities from last instar larvae of *Z. trifolii* (lower panel). ep: epidermis; cav I: type I cuticular cavity; cav II: type II cuticular cavity. Adapted from (Naumann et al., 1999).

identical to the pathway in plants (Conn, 1991; Davis and Nahrstedt, 1987; Holzkamp and Nahrstedt, 1994; Nahrstedt, 1996; Wray et al., 1983) (Fig. 1).

$\beta$ -Glucosidase dependant HCN release has been observed from different life stages of many zygaenid species (Davis and Nahrstedt, 1979; Witthohn and Naumann, 1987b).  $\beta$ -Glucosidase from *Z. trifolii* is a dimer consisting of two supposedly identical 66 kDa subunits. The  $\beta$ -glucosidase exhibits a strong activity towards the endogenous substrates linamarin and lotaustralin, with lotaustralin being a better substrate (Franzl et al., 1989). The *Z. trifolii*  $\beta$ -glucosidase is labile at temperatures above 40 °C and inactivated at a pH below 5 as are  $\beta$ -glucosidases from other insects (*Bombyx mori* and *Trinervitermes trinervoides*).  $\beta$ -Glucosidase activity was found exclusively in haemolymph, which has a pH of 6.2 (Franzl et al., 1989). At pH 6.2, the  $\beta$ -glucosidase is present in an almost inactive state but the enzyme becomes active when pH decreases (Nahrstedt and Müller, 1993). This may point to a situation where stomach acid from a predator will activate the  $\beta$ -glucosidase leading to a rapid and strong release of HCN (Nahrstedt, 1993).  $Mg^{++}$  and  $Ca^{++}$  ions are inhibitors of the *Zygaena*  $\beta$ -glucosidase. To date, this is the only example of a  $\beta$ -glucosidase which is inhibited by alkaline earth metal ions (Nahrstedt and Müller, 1993).

$\alpha$ -Hydroxynitrile lyase was purified from the haemolymph of *Z. trifolii* and characterized as a FAD-containing enzyme (Müller and Nahrstedt, 1990; Nahrstedt, 1996).

$\beta$ -Cyanoalanine synthase activity is found in the integument (22%), the fat body (27%) and the gut (12%) of *Zygaena* larvae, with the highest activity of the enzyme in the gut.  $\beta$ -Cyanoalanine synthase detoxifies HCN to  $\beta$ -cyanoalanine, which accumulates in haemolymph (75%) and integument (16%) (Nahrstedt, 1993; Witthohn and Naumann, 1987a). Thus, these insects can easily detoxify HCN. Zygaenidae seem to use the  $\beta$ -cyanoalanine pathway as the sole pathway for detoxification of HCN, since rhodanese (Fig. 1) has not been found in any species containing  $\beta$ -cyanoalanine (Witthohn and Naumann, 1987a).

## 6. Cyanogenic glucosides in Papilionoidea (butterflies)

### 6.1. Linamarin and lotaustralin distribution

Imagines of butterflies from the Heliconiinae, Acraeinae, Nymphalinae and Polyommatae (Papilionoidea) groups (Fig. 3) accumulate linamarin and often lotaustralin in all life stages (Brown and Francini, 1990; Nahrstedt, 1988; Nahrstedt and Davis, 1981; 1983). The butterflies *de novo* synthesize linamarin and lotaustralin from valine and isoleucine, respectively, and because

linamarin and lotaustralin are not present in their larval host plants (Passifloraceae), sequestration cannot occur (Engler et al., 2000; Nahrstedt and Davis, 1983; Wray et al., 1983). The amount of linamarin is higher than that of lotaustralin as also observed in imagines of Zygaenidae, and the CNGs have the same bodily distribution as observed in *Z. trifolii* (Davis and Nahrstedt, 1982; Franzl et al., 1986). The examined butterflies also contain monoglycoside cyclopentenyl cyanogens, probably sequestered from their host plants, at the larval stage (Engler et al., 2000).

### 6.2. Detoxification, biosynthesis and sequestration of cyanogenic glucosides

In *Heliconius sara* (Papilionoidea) (Fig. 3), monoglycoside cyclopentenyl cyanogens, obtained and sequestered from the host plant are detoxified using a unique enzymatic mechanism not found in the host plants. This mechanism involves substitution of the nitrile group of the cyanohydrin function with a mercapto group (Engler et al., 2000). The reaction mechanism involved has not been elucidated and it remains to be shown whether free cyanide is released during the reaction or whether the nitrile group is transferred to a specific acceptor molecule, like cysteine, as a part of the reaction sequence. The cyanogenic glucoside-derived mercapto compound was not found in the host plant (Engler et al., 2000). This is in contrast to the mechanism proposed for *H. melpomene* (Papilionoidea) (Fig. 3) and *Z. trifolii*. The presence of  $\beta$ -cyanoalanine synthase was demonstrated in *Clossiana euphrosyne* (Papilionoidea) (Witthohn and Naumann, 1987a) (Fig. 3) which confirms that some *Papilionoidea* species detoxify HCN using the same mechanism as Zygaenidae species. Consequently, *H. sara* may either have a unique detoxification system, or insects may use different detoxification systems within the same group.

*Eutoptia hegesia* (Heliconiinae, Papilionoidea) (Fig. 3) has been hypothesized to *de novo* synthesize as well as sequester cyclopentenyl glycine derived CNGs from its primary host plant *Turnera ulmifolia* (Schappert and Shore, 1999). *De novo* biosynthesis was proposed because HCN was measured in larvae reared on acyanogenic plants. Sequestering was proposed because of significantly higher CNG levels in larvae reared on cyanogenic plants compared to siblings on acyanogenic plants (Schappert and Shore, 1999). Accordingly, the ability to both *de novo* synthesize and sequester the same CNGs appears not to be an exclusive feature for Zygaenidae species.

Within Papilionoidea, *de novo* biosynthesis of CNGs seem to be less frequent in adults of less advanced genera such as *Agraulus* (Heliconiinae, Papilionoidea), *Dryas* (Heliconiinae, Papilionoidea) and *Cethosia*

(Acraeinae, Papilionoidea) (Fig. 3) compared to the more advanced genera *Heliconius* (Nahrstedt and Davis, 1985). In all Heliconiini (Papilionoidea) and some *Acraea* (Papilionoidea) species (Fig. 3), the ability to synthesize and metabolize CNGs as well as to detoxify HCN may have provided the possibility to change from acyanogenic host plants to the cyanogenic Passifloraceae (Davis and Nahrstedt, 1985). Synthesis and degradation of CNGs in all these butterfly species may have been derived from a common cyanogenic ditrysian ancestor (Fig. 3) and may therefore be homologous to the pathways found in Zygaenidae.

## 7. Conclusions and perspectives

In plants, cyanogenic glucosides serve as good chemotaxonomic markers for plant relatedness: the more closely related two plant species are, the more similar their cyanogenic glucosides are (Jones, 1988). Within cyanogenic genera, however, acyanogenic species can be found. These are the result of natural mutations that have caused the loss of the ability to produce one or more of the enzymes involved in cyanogenic glucoside biosynthesis and/or in the degradation of CNGs. Secondary loss of enzymes involved in biosynthesis or degradation of CNGs in insects may also be found upon further study. Eventually, it may become apparent that biosynthesis and degradation of CNGs and the ability to detoxify HCN are shared traits between groups of moths and butterflies, and derived from a common cyanogenic ditrysian ancestor (Fig. 3). Studies on a range of ditrysian species are required before this hypothesis can be substantiated or dismissed.

Acyanogenic *L. corniculatus* plants are grazed more heavily by herbivores than cyanogenic *L. corniculatus* plants, but the difference is partly balanced by the ability of acyanogenic plants to grow better than cyanogenic ones (Jones, 1962, 1988). Accordingly, it is probably advantageous for *L. corniculatus* to maintain this polymorphism. This would serve to secure more competitive growth and development in years with no or reduced numbers of herbivores, and to prevail in years with heavy attack from herbivores. Acquisition of an ability to detoxify HCN in some insect species early in their evolution, has offered the opportunity to feed on plants that deter other herbivores, and subsequently exploit the availability of such new host plants to radiate at the expense of otherwise competing species, as seen in moth and butterfly groups. Subsequent acquisition of the ability to sequester CNGs from host plants as a means of detoxification provides such insects with the additional benefits of an improved defense system. Simultaneously, these insects then become more or less dependent on the availability of cyanogenic host

plants. In this scenario, it is advantageous for the plant to maintain some acyanogenic genotypes that will not be preferred by the specialized insects. The ability to de novo synthesize CNGs is probably a basic trait within some insect groups. Accordingly, these insects only need cyanogenic host plants to minimize their own biosynthesis of CNGs. Consequently, some moth and butterfly species presumably feed on plants that deter other herbivores without being absolutely dependant on such plants for their own predator defense.

The ability to transfer genes across species using genetic engineering enables the design of plants with an altered qualitative and quantitative content of natural products thereby bypassing millions of years of co-evolution of plants and their herbivores. Transgenic *A. thaliana* plants accumulating the tyrosine-derived CNG dhurrin provided strong evidence that dhurrin serves as a strong feeding deterrent in free choice experiments with the flea beetle *Phyllotreta nemorum* (Tattersall et al., 2001). Similarly, transgenic *Lotus japonicus* plants with altered ratios of linamarin and lotaustralin are available (Forslund et al., submitted) to study sequestering and turnover of linamarin and lotaustralin in *Z. transalpiina*. In this way, chemical warfare between plants and insects can be followed closely through metabolite profiling (LC-MS) and transcript profiling, thereby providing a detailed understanding of the relative importance of complete metabolism, detoxification and sequestering of CNGs.

Many details remain unknown concerning synthesis and degradation of CNGs in insects. Has the ability to de novo synthesize CNGs evolved several times in the course of insect evolution? Or have genes encoding the enzymes involved in biosynthesis, degradation and detoxification of CNGs been horizontally transferred to the insects from e.g. host plants? If the genes have been transferred, did this then happen early in insect evolution or on several occasions? Or is the ability of some insects to synthesize, degrade and detoxify CNGs the result of convergent evolution, where modifications of distantly related genes have enabled recruitment of new desired functions? As more insect species become analyzed and additional information is obtained, answers to some of these questions will be provided. It should be noticed that already with the sparse knowledge currently available, it is obvious that some of the plant enzymes involved in biosynthesis or degradation of cyanogenic glucosides, like the  $\alpha$ -hydroxynitrilases, have been recruited by modification of different types of ancestral genes. This may also apply to the arthropods. Valuable new information will appear as more insect genomes, e.g. *Manduca sexta*, *Heliothis virescens*, and *Heliconius erato* (National Human Genome Research Institute), are sequenced.

## Acknowledgements

We thank Professors Peter Esbjerg and Jørgen Eilenberg as well as Curator Jan Martin from Department of Ecology for help with handling *Zygaena* insects. We are grateful to Dr. Carl Erik Olsen for LC-MS analyses. This work was supported by the Danish National Research Foundation and the Danish Veterinary & Agricultural Research council.

## References

- Andersen, M.D., Busk, P.K., Svendsen, I., Møller, B.L., 2000. Cytochromes P-450 from cassava (*Manihot esculenta* Crantz) catalyzing the first steps in the biosynthesis of the cyanogenic glucosides linamarin and lotaustralin. *J. Biol. Chem.* 275, 1966–1976.
- Bak, S., Kahn, R.A., Nielsen, H.L., Møller, B.L., Halkier, B.A., 1998. Cloning of three A type cytochromes P450, CYP71E1, CYP98, and CYP99 from *Sorghum bicolor* (L.) Moench by a PCR approach and identification by expression in *Escherichia coli* of CYP71E1 as a multifunctional cytochrome P450 in the biosynthesis of the cyanogenic glucoside dhurrin. *Plant Mol. Biol.* 36, 393–405.
- Beesley, S.G., Compton, S.G., Jones, D.A., 1985. Rhodanese in insects. *J. Chem. Ecol.* 11, 45–50.
- Bernays, E.A., 1991. Relationship between deterrence and toxicity of plant secondary compounds for the grasshopper *Schistocerca americana*. *J. Chem. Ecol.* 17, 2519–2526.
- Bernays, E.A., Cornelius, M., 1992. Relationship between deterrence and toxicity of plant secondary compounds for the alfalfa weevil *Hypera brunneipennis*. *Entomologia Experimentalis et Applicata* 64, 289–292.
- Bordo, D., Bork, P., 2002. The rhodanese/Cdc25 phosphatase superfamily—sequence–structure–function relations. *EMBO Reports* 3, 741–746.
- Braekman, J.C., Daloz, D., Pasteels, J.M., 1982. Cyanogenic and other glucosides in a Neo-Guinean bug *Leptocoris isolata*—possible precursors in its host-plant. *Biochem. Syst. Ecol.* 10, 355–364.
- Brattsten, L.B., Samuelian, J.H., Long, K.Y., Kincaid, S.A., Evans, C.K., 1983. Cyanide as a feeding stimulant for the southern armyworm, *Spodoptera eridania*. *Ecol. Entomology* 8, 125–132.
- Brown, K.S., Francini, R.B., 1990. Evolutionary strategies of chemical defense in aposematic butterflies: cyanogenesis in Asteraceae-feeding American Acreinae. *Chemoeology* 1, 52–56.
- Catric, P., Farnden, K., Conn, E.E., 1972. Cyanide metabolism in higher plants: V. The formation of asparagine from  $\beta$ -cyanoalanine. *Arch. Biochem. Biophys.* 152, 62–69.
- Cicek, M., Blanchard, D.J., Bevan, D.R., Esen, A., 2000. The aglycone specificity-determining sites are different in 2,4-dihydroxy-7-methoxy-1,4-benzoxazin-3-one (DIMBOA)-glucosidase (maize  $\beta$ -glucosidase) and dhurrinase (*Sorghum*  $\beta$ -glucosidase). *J. Biol. Chem.* 275, 20002–20011.
- Cicek, M., Esen, A., 1998. Structure and expression of a dhurrinase ( $\beta$ -glucosidase) from *Sorghum*. *Plant Physiol.* 116, 1469–1478.
- Compton, S.G., Jones, D.A., 1985. An investigation of the responses of herbivores to cyanogenesis in *Lotus corniculatus*. *Biol. J. Linn. Soc.* 26, 21–38.
- Conn, E.E., 1980. Cyanogenic compounds. *Ann. Rev. Plant Physiol.* 31, 433–451.
- Conn, E.E., 1991. The metabolism of a natural product—lessons learned from cyanogenic glycosides. *Planta Medica* 57, S1–S9.
- Czjzek, M., Cicek, M., Zamboni, V., Bevan, D.R., Henrissat, B., Esen, A., 2000. The mechanism of substrate (aglycone) specificity in  $\beta$ -glucosidases is revealed by crystal structures of mutant maize  $\beta$ -glucosidase-DIMBOAGlc, and -dhurrin complexes. *Proc. Natl. Acad. Sci. U.S.A.* 97, 13555–13560.
- Davis, R.H., Nahrstedt, A., 1979. Linamarin and lotaustralin as the source of cyanide in *Zygaena filipendulae* (Lepidoptera). *Comp. Biochem. Physiol. B-Biochemistry & Molecular Biology* 64, 395–397.
- Davis, R.H., Nahrstedt, A., 1982. Occurrence and variation of the cyanogenic glucosides linamarin and lotaustralin in species of the Zygaenidae (Insecta, Lepidoptera). *Comp. Biochem. Physiol. B-Biochemistry & Molecular Biology* 71, 329–332.
- Davis, R.H., Nahrstedt, A., 1985. Cyanogenesis in insects. In: Kerkut, G.A., Gilbert, L.I. (Eds.), *Comprehensive Insect Physiology, Biochemistry and Pharmacology*. Pergamon Press, Oxford, pp. 635–654.
- Davis, R.H., Nahrstedt, A., 1987. Biosynthesis of cyanogenic glucosides in butterflies and moths—effective incorporation of 2-methylpropanenitrile and 2-methylbutanenitrile into linamarin and lotaustralin by *Zygaena* and *Heliconius* species (Lepidoptera). *Insect Biochem.* 17, 689–693.
- Duffey, S.S., 1981. Cyanide and arthropods. In: Vennesland, B., Conn, E.E., Knowles, C.J., Westley, J., Wissing, F. (Eds.), *Cyanide in Biology*. Academic Press, London, pp. 385–414.
- Duffey, S.S., Towers, G.H.N., 1978. On the biochemical basis of HCN production in the millipede *Harpaghe haydeniana* (Xystodesmidae: Polydesmida). *Can. J. Zool.* 56, 7–16.
- Engler, H.S., Spencer, K.C., Gilbert, L.E., 2000. Insect metabolism—preventing cyanide release from leaves. *Nature* 406, 144–145.
- Esen, A., 1993.  $\beta$ -Glucosidases—overview. *ACS. Symp. Series* 533, 1–14.
- Feyereisen, R., 1999. Insect P450 enzymes. *Ann. Rev. Entomology* 44, 507–533.
- Franzl, S., Ackermann, I., Nahrstedt, A., 1989. Purification and characterization of a  $\beta$ -glucosidase (linamarase) from the haemolymph of *Zygaena trifolii* Esper, 1783 (Insecta, Lepidoptera). *Experientia* 45, 712–718.
- Franzl, S., Nahrstedt, A., Naumann, C.M., 1986. Evidence for site of biosynthesis and transport of the cyanoglucosides linamarin and lotaustralin in larvae of *Zygaena trifolii* (Insecta, Lepidoptera). *J. Insect Physiol.* 32, 705–709.
- Franzl, S., Naumann, C.M., 1985. Cuticular cavities—storage chambers for cyanoglucoside—containing defensive secretions in larvae of a Zygaenid moth. *Tissue & Cell* 17, 267–278.
- Franzl, S., Naumann, C.M., Nahrstedt, A., 1988. Cyanoglucoside storing cuticle of *Zygaena* larvae (Insecta, Lepidoptera)—morphological and cyanoglucoside changes during the molt. *Zoomorphology* 108, 183–190.
- Gaunt, M.W., Miles, M.A., 2002. An insect molecular clock dates the origin of the insects and accords with paleontological and biogeographic landmarks. *Mol. Biol. Evol.* 19, 748–761.
- Gebrehiwot, L., Beuselinck, P.R., 2001. Seasonal variations in hydrogen cyanide concentration of three *Lotus* species. *Agronomy J.* 93, 603–608.
- Gleadow, R.M., Woodrow, I.E., 2002. Constraints on effectiveness of cyanogenic glycosides in herbivore defense. *J. Chem. Ecol.* 28, 1301–1313.
- Halkier, B.A., 1996. Catalytic reactivities and structure/function relationships of cytochrome P450 enzymes. *Phytochemistry* 43, 1–21.
- Halkier, B.A., Møller, B.L., 1989. Biosynthesis of the cyanogenic glucoside dhurrin in seedlings of *Sorghum bicolor* (L.) Moench and partial purification of the enzyme system involved. *Plant Physiol.* 90, 1552–1559.
- Hansen, K.S., Kristensen, C., Tattersall, D.B., Jones, P.R., Olsen, C.E., Bak, S., 2003. Møller, B. L., The in vitro substrate regioselectivity of UGT85B1, the cyanohydrin glucosyltransferase from *Sorghum bicolor*. *Phytochemistry* 64, 143–151.

- Hickel, A., Hasslacher, M., Griengl, H., 1996. Hydroxynitrile lyases: functions and properties. *Physiologia Plantarum* 98, 891–898.
- Holzkamp, G., Nahrstedt, A., 1994. Biosynthesis of cyanogenic glucosides in the Lepidoptera—incorporation of [U-C-14]-2-methylpropanealdoxime, 2S-[U-C-14]-methylbutanealdoxime and D,L-[U-C-14]-N-hydroxyisoleucine into linamarin and lotaustralin by the larvae of *Zygaena trifolii*. *Insect Biochem. Mol. Biol.* 24, 161–165.
- Hösel, W., Conn, E.E., 1982. The aglycone specificity of plant  $\beta$ -glucosidases. *Trends Biochem. Sci.* 7, 219–221.
- Hösel, W., Tober, I., Eklund, S.H., Conn, E.E., 1987. Characterization of  $\beta$ -glucosidases with high specificity for the cyanogenic glucoside dhurrin in *Sorghum bicolor* (L.) Moench seedlings. *Arch. Biochem. Biophys.* 252, 152–162.
- Hu, Z., Poulton, J.E., 1997. Sequencing, genomic organization, and preliminary promoter analyses of a black cherry (R)(+)-mandelonitrile lyase gene. *Plant Physiol.* 115, 1359–1369.
- Hu, Z., Poulton, J.E., 1999. Molecular analysis of (R)(+)-mandelonitrile lyase microheterogeneity in black cherry. *Plant Physiol.* 11, 1535–1546.
- Ikegami, F., Murakoshi, I., 1994. Enzyme synthesis of non-protein  $\beta$ -substituted alanines and some higher homologues in plants. *Phytochemistry* 35, 1089–1104.
- Ikegami, F., Takayama, K., Murakoshi, I., 1988. Purification and properties of  $\beta$ -cyanoalanine synthase from *Lathyrus latifolius*. *Phytochemistry* 27, 3385–3389.
- Jones, D.A., 1988. Cyanogenesis in animal–plant interactions. *Ciba. Found. Symp.* 140, 151–170.
- Jones, D.A., Rothschild, M., Parsons, J., 1962. Release of hydrocyanic acid from crushed tissues of all stages in life-cycle of species of Zygaeninae (Lepidoptera). *Nature* 193, 52.
- Jones, D.A., 1962. Selective eating of the acyanogenic form of the plant *Lotus corniculatus* by various animals. *Nature* 4820, 1109–1110.
- Jones, P.R., Andersen, M.D., Nielsen, J.S., Høj, P.B., Möller, B.L., 2000. The biosynthesis, degradation, transport and possible function of cyanogenic glucosides. In: Romero, J.T., Ibrahim, R., Varin, L., De Luca, V. (Eds.), *Evolution of Metabolic Pathways*. Elsevier Science, New York, pp. 191–247.
- Jones, P.R., Möller, B.L., Høj, P.B., 1999. The UDP-glucose : *p*-hydroxymandelonitrile-*O*-glucosyltransferase that catalyzes the last step in synthesis of the cyanogenic glucoside dhurrin in *Sorghum bicolor*—isolation, cloning, heterologous expression, and substrate specificity. *J. Biol. Chem.* 274, 35483–35491.
- Kahn, R.A., Bak, S., Svendsen, I., Halkier, B.A., Möller, B.L., 1997. Isolation and reconstitution of cytochrome P450ox and in vitro reconstitution of the entire biosynthetic pathway of the cyanogenic glucoside dhurrin from sorghum. *Plant Physiol.* 115, 1661–1670.
- Lechtenberg, M., Nahrstedt, A., 1999. Cyanogenic glucosides. In: Ikan, R. (Ed.), *Naturally Occurring Glycosides*. John Wiley & Sons, New York, pp. 147–191.
- Levinson, N.Z., Kaissling, K.-E., Levinson, A.R., 1973. Olfaction and cyanide sensitivity in the six-spot burnet moth *Zygaena fillipendulae* and the silk moth *Bombyx mori*. *J. Comp. Physiol.* 86, 209–215.
- Meyers, D., Ahmad, S., 1991. Link between L-3-cyanoalanine synthase activity and differential cyanide sensitivity of insects. *Biochim. Biophys. Acta* 1075, 195–197.
- Miller, J.M., Conn, E.E., 1980. Metabolism of hydrogen cyanide by higher plants. *Plant Physiol.* 65, 1199–1202.
- Möller, B.L., Conn, E.E., 1980. The biosynthesis of cyanogenic glucosides in higher plants. Channeling of intermediates in dhurrin biosynthesis by a microsomal system from *Sorghum bicolor* (L.). *Moench. J. Biol. Chem.* 255, 3049–3056.
- Möller, B.L., Poulton, J.E., 1993. Cyanogenic glucosides. In: Lea, P.J. (Ed.), *Methods in Plant Biochemistry*, vol. 9. Academic Press, San Diego, pp. 183–207.
- Möller, B.L., Seigler, D.S., 1999. Biosynthesis of cyanogenic glucosides, cyanolipids and related compounds. In: Singh, B.K. (Ed.), *Plant Amino Acids*. M. Dekker, New York, pp. 563–609.
- Morant, M., Bak, S., Möller, B.L., Werck-Reichhart, D., 2003. Plant cytochromes P450: tools for pharmacology, plant protection and phytomediation. *Curr. Op. Biotechnology* 14, 151–162.
- Müller, E., Nahrstedt, A., 1990. Purification and characterization of an  $\alpha$ -hydroxynitrile lyase from the haemolymph of the larvae of *Zygaena trifolii*. *Planta Medica* 56, 611–612.
- Müller, T., Ockenfels, P., Naumann, C.M., 1993. Olfactory discrimination between host- and non-host-plants in *Zygaena trifolii* (Esper, 1783) and *Z. transalpiina* (Esper, 1780) (Lepidoptera, Zygaenidae). In: Tremewan, W.G., Wipking, W., Naumann, C.M. (Eds.), *Proceedings of the 5th International Symposium on the Biology of the Zygaenidae (Insecta, Lepidoptera)*. Koeltz Scientific Books, Grietherbusch (Germany), pp. 125–137.
- Nahrstedt, A., 1985. Cyanogenic compounds as protecting agents for organisms. *Plant Syst. Evol* 150, 35–47.
- Nahrstedt, A., 1988. Cyanogenesis and the role of cyanogenic compounds in insects. *Ciba. Found. Symp* 140, 131–150.
- Nahrstedt, A., 1989. The significance of secondary metabolites for interactions between plants and insects. *Planta Medica* 333–338.
- Nahrstedt, A., 1993. Cyanogenesis in the Zygaenidae (Lepidoptera): a review of the state of the art. In: Tremewan, W.G., Wipking, W., Naumann, C.M. (Eds.), *Proceedings of the 5th International Symposium on the Biology of the Zygaenidae (Insecta, Lepidoptera)*. Koeltz Scientific Books, Grietherbusch (Germany), pp. 17–29.
- Nahrstedt, A., 1996. Relationships between the defensive systems of plants and insects. In: Romeo, X., Saunders, X., Barbossa, X. (Eds.), *Recent Advances in Phytochemistry*. Plenum Press, New York, pp. 217–230.
- Nahrstedt, A., Davis, R.H., 1981. The occurrence of the cyanoglucosides, linamarin and lotaustralin, in *Acræa* and *Heliconius* butterflies. *Comp. Biochem. Physiol. B—Biochemistry & Molecular Biology* 68, 575–577.
- Nahrstedt, A., Davis, R.H., 1983. Occurrence, variation and biosynthesis of the cyanogenic glucosides linamarin and lotaustralin in species of the Heliconiini (Insecta, Lepidoptera). *Comp. Biochem. Physiol. B—Biochemistry & Molecular Biology* 75, 65–73.
- Nahrstedt, A., Davis, R.H., 1985. Biosynthesis and quantitative relationships of the cyanogenic glucosides, linamarin and lotaustralin, in genera of the Heliconiini (Insecta, Lepidoptera). *Comp. Biochem. Physiol. B—Biochemistry & Molecular Biology* 82, 745–749.
- Nahrstedt, A., Müller, E., 1993.  $\beta$ -Glucosidase (linamarase) of the larvae of the moth *Zygaena trifolii* and its inhibition by some alkaline-earth metal-ions. *ACS. Symp. Series* 533, 132–144.
- Naumann, C.M., Tarmann, G.M., Tremewan, W.G., 1999. *The Western Palaearctic Zygaenidae: (Lepidoptera)*. Apollo Books, Stenstrup, Denmark.
- Paquette, S.M., Möller, B.L., Bak, S., 2003. On the origin of family 1 plant glycosyltransferases. *Phytochemistry* 62, 399–413.
- Poulton, J.E., 1990. Cyanogenesis in plants. *Plant Physiol.* 94, 401–405.
- Ranson, H., Claudianos, C., Ortelli, F., Abgrall, C., Hemingway, J., Sharakhova, M.V., Unger, M.F., Collins, F.H., Feyereisen, R., 2002. Evolution of supergene families associated with insecticide resistance. *Science* 298, 179–181.
- Ressler, C., Nigam, S., Giza, Y., 1969. Toxic principle in vetch: isolation and identification of  $\gamma$ -L-glutamyl-L- $\beta$ -cyanoalanine from common vetch seeds: distribution in some legumes. *J. Amer. Chem. Soc.* 91, 2758–2765.
- Schappert, P.J., Shore, J.S., 1999. Effects of cyanogenesis polymorphism in *Turnera ulmifolia* on *Euptoieta hegesia* and potential *Anolis* predators. *J. Chem. Ecol.* 25, 1455–1479.
- Sibbesen, O., Koch, B., Halkier, B.A., 1994. Möller, B. L., Isolation of the heme-thiolate enzyme cytochrome P450Tyr, which catalyzes the committed step in the biosynthesis of the cyanogenic glucoside dhurrin in *Sorghum bicolor* (L.) Moench. *Proc. Natl. Acad. Sci. USA* 91, 9740–9744.

- Swain, E., Li, C.P., Poulton, J.E., 1992. Tissue and subcellular localization of enzymes catabolizing (*R*)-amygdalin in mature *Prunus serotina* seeds. *Plant Physiol.* 100, 291–300.
- Tattersall, D.B., Bak, S., Jones, P.R., Olsen, C.E., Nielsen, J.K., Hansen, M.L., Høj, P.B., Møller, B.L., 2001. Resistance to an herbivore through engineered cyanogenic glucoside synthesis. *Science* 293, 1826–1828.
- Tattersall, D.B., Nielsen, K.A., Jones, P.R., Møller, B.L., 2003. Metabolon formation in dhurrin biosynthesis as documented by confocal laser scanning microscopy. *Plant Cell* (in press).
- Tijet, N., Helvig, C., Feyereisen, R., 2001. The cytochrome P450 gene superfamily in *Drosophila melanogaster*: annotation, intron–exon organization and phylogeny. *Gene* 262, 189–198.
- Trummel, K., Wajant, H., 1997. Molecular cloning of acetone cyanohydrin lyase from flax (*Linum usitatissimum*). Definition of a novel class of hydroxynitrile lyases. *J. Biol. Chem.* 272, 4770–4774.
- Vetter, J., 2000. Plant cyanogenic glycosides. *Toxicol.* 38, 11–36.
- Vogt, T., Jones, P., 2000. Glycosyltransferases in plant natural product synthesis: characterization of a supergene family. *Trends Plant Sci.* 5, 380–386.
- Wajant, H., Pfizenmaier, K., 1996. Identification of potential active-site residues in the hydroxynitrile lyase from *Manihot esculenta* by site-directed mutagenesis. *J. Biol. Chem.* 271, 25830–25834.
- Werck-Reichhart, D., Bak, S., Paquette, S.M., 2002. Cytochromes P450. In: Somerville, C.R., Meyerowitz, E.M. (Eds.), *The Arabidopsis Book*. American Society of Plant Biologists, Rockville, pp. 1–28. (tab.0028, <http://www.aspb.org/publications/arabidopsis/>).
- Werck-Reichhart, D., Feyereisen, R., 2000. Cytochromes P450: a success story. *Genome Biology* 1, 3003.1–3003.9.
- Werck-Reichhart, D., Hehn, A., Didierjean, L., 2000. Cytochromes P450 for engineering herbicide tolerance. *Trends Plant Sci.* 5, 116–123.
- Witthohn, K., Naumann, C.M., 1984. Qualitative and quantitative studies on the compounds of the larval defensive secretion of *Zygaena trifolii* (Esper, 1783) (Insecta, Lepidoptera, Zygaenidae). *Comp. Biochem. Physiol. C—Pharmacology Toxicology & Endocrinology* 79, 103–106.
- Witthohn, K., Naumann, C.M., 1987a. Cyanogenesis—a general phenomenon in the Lepidoptera. *J. Chem. Ecol.* 13, 1789–1809.
- Witthohn, K., Naumann, C.M., 1987b. Genus *Zygaena* F and related taxa (Insecta, Lepidoptera). 53. Active cyanogenesis—in Zygaenidae and other Lepidoptera. *Z. Naturforsch* 42c, 1319–1322.
- Wray, V., Davis, R.H., Nahrstedt, A., 1983. Biosynthesis of cyanogenic glycosides in butterflies and moths—incorporation of valine and isoleucine into linamarin and lotaustralin by *Zygaena* and *Heliconius* species (Lepidoptera). *Z. Naturforsch* 38c, 583–588.
- Yip, W.-K., Yang, S., 1988. Cyanide metabolism in relation to ethylene production in plant tissues. *Plant Physiol.* 88, 473–476.

with special focus on biosynthesis, sequestration and degradation of linamarin and lotaustralin in *Zygaena transalpiina* and characterization of the enzyme systems and genes involved.



**Søren Bak** obtained his MSc in Biochemistry at the University of Copenhagen in 1993 and his PhD in Plant Molecular Biology at the Royal Veterinary & Agricultural University in 1997. In his PhD Thesis, new cytochrome P450s in primary and secondary metabolism were identified and characterized in *Sorghum bicolor*. As a Post Doctoral Fellow at the Plant Biochemistry Laboratory, Department of Plant Biology, he initiated work on metabolic engineering of natural products in *Arabidopsis thaliana*. This was followed by a two year stay at Department of Plant Sciences, University of Arizona where he worked on functional genomics of the cytochrome P450 multigene family in *A. thaliana*. Since 2001 he has been employed as Associate Professor at the Department of Plant Biology, The Royal Veterinary & Agricultural University, Copenhagen. His work is focused on biochemical, molecular and evolutionary aspects of natural products with a specific emphasis on cyanogenic glucosides, cytochromes P450, and family I glycosyltransferases. He hosts the bioinformatics site *The Arabidopsis P450, Cytochromes b5, P450 Reductase, and Glycosyltransferase Site at PlaCe*, (<http://www.biobase.dk/P450>).



**Anne Vinther Rasmussen** obtained her MSc in Biology at the University of Copenhagen in 2003. For her MSc Thesis, she identified, cloned and heterologously expressed new genes encoding  $\beta$ -ketoacyl [acyl carrier protein] synthases. She began her PhD studies at the Plant Biochemistry Laboratory, Royal Veterinary & Agricultural University, Copenhagen in March 2003 with the aim of developing *Lotus japonicus* as a genetic model system for studies on the function of cyanogenic glucosides in plants and their role in plant–insect interactions.

A key approach is use of metabolic engineering to block synthesis of, or alter, the composition of cyanogenic glucosides in this cyanogenic legume.



**Mika Zagrobelny** obtained her MSc in Biology at the University of Copenhagen in 2002. For her MSc Thesis she conducted an exhaustive study of the Sam/Frodo family of non-LTR (Long Terminal Repeat) retrotransposons in the genomes of the two nematodes *Caenorhabditis elegans* and *C. briggsae* with special focus on the function of transposable elements as integrated parts of the genome. She is currently employed as a Research Assistant at the Center of Molecular Plant Physiology (PlaCe) at the Royal Veterinary & Agricultural University, Copenhagen, where she works with cyanogenic glucosides in insects



**Bodil Jørgensen** obtained her MSc in Biology at the University of Copenhagen in 1989. She carried out her PhD studies on genetic transformation and regeneration of pea at the plant breeding company Maribo Seeds, a subsidiary of Danisco A/S and obtained her PhD from the University of Copenhagen in 1992. She spent three years as a Post Doctoral Fellow at the Institute of Grassland and Environmental Research, Aberystwyth, Wales where the focus of her work was identification and characterization of genes responsible for root nodulation and nitrogen fixation. Since 1997,

she has been employed as Senior Scientist at the Danish Institute of Agricultural Science (DIAS). Her current research focus is to develop transformation systems for marker free GMO crops that are vegetatively propagated.



**Clas M. Naumann** obtained his PhD at the University of Bonn in 1970. He was employed as Associate Professor at the universities of Kabul (Afghanistan) and Bonn until 1974 when he moved to Munich. In 1978, he was appointed Professor in Zoology at the University of Bielefeld but returned to Bonn to fill the Chair as Professor of Zoology and Director of the Zoological Science Institute (Zoologisches Forschungsinstitut) and Alexander König Museum. Among a number of trusted positions, he served as an elected referee

of Deutsche Forschungsgemeinschaft (DFG) from 1992 to 2000. He is Chief Editor of the *Zoologischer Anzeiger*, Editor of Handbook of Palaearctic Macrolepidoptera and *Entomologische Zeitschrift*, but also serves other journals, such as *Fauna of Saudi Arabia*, *Zoology in the Middle East*, *Acta Zoologica Hungarica* and *Entomologica Germanica*. His main research interests are within the field of insect phylogeny and taxonomy with a special focus on Lepidoptera. Other major research contributions are within evolutionary biology, biogeography, evolution and ecology.



**Birger Lindberg Møller** obtained his MSc, PhD and DSc from the University of Copenhagen in 1972, 1975 and 1984, respectively. His MSc Thesis work was focussed on identification of alkaloids in African medicinal plants. The topic of his PhD Thesis was lysine metabolism in plants and the experimental work was carried out at the Royal Veterinary & Agricultural University under the supervision of Peder Olesen Larsen. He spent the next three years as a Fulbright Fellow at Eric Conn's laboratory, University of California, Davis, working with biosynthesis of cyanogenic glucosides.

From 1977 to 1984 he was Senior Scientist and Niels Bohr Fellow at the Department of Physiology, Carlsberg Laboratory with Diter von Wettstein and studying photosynthesis. This work formed the basis for his DSc Thesis. In 1984 he was appointed Research Professor and in 1989 Professor in Plant Biochemistry at the Royal Veterinary & Agricultural University, Copenhagen. In 1998 he became the Head of Center for Molecular Plant Physiology (PlaCe). One of his main research interests within plant biochemistry is synthesis, storage and degradation of cyanogenic glucosides and elucidation of their role in plant insect and plant microbe interactions. The knowledge obtained is being used to ameliorate important crop plants like barley and cassava with respect to nutritive value and pest resistance. He is member of the Senate of the Royal Veterinary & Agricultural University, Vice-Chairman of its Research Committee, Board member of the Danish Institute of Agricultural Science and member of the Board of Trustees of the International Institute of Tropical Agriculture (IITA) in Ibadan, Nigeria. He is a co-founder of the biotech company Poalis.